



LEVERAGING TECHNOLOGY FOR BETTER OUTCOMES

Improving Lives of Patients & Clinicians





SPEAKER DISCLOSURE

I have nothing to disclose.

Neuraxial Anesthesia for Vaginal Delivery in Congenital Complete Heart Block (CCHB)

Background:


- **Congenital Heart Block Incidence:** ~1 in 20,000 pregnancies [1]
- Many remain asymptomatic due to adequate **chronotropic compensation** [1]
- Fluoroscopy needed for pacemaker insertion poses **fetal risks**, making intervention decisions complex.
- **Neuraxial anesthesia:** Can reduce sympathetic tone → unmask/exacerbate bradyarrhythmia [2]
- **Baseline HR <60 bpm:** High risk for severe peripartum bradycardia [2]
- **Aim:** Present anesthetic and obstetric management of a parturient with CCHB



[1] Sundararaman, L., Cohn, J. H., & Ranasinghe, J. S. (2016). Complete heart block in pregnancy: case report, analysis, and review of anesthetic management. *Journal of clinical anesthesia*, 33, 58-61. p

Case:

- **Patient Profile:** 26-year-old, gravida 2 para 1, at 35.2 weeks gestation with known CCHB.
- **Clinical Status:**
 - HR: 51–59 bpm, BP: 130–140/70–80 mmHg, SpO₂ >98%
 - Known CCHB → Early induction for **worsening conduction** concerns
 - **Multidisciplinary consultation:** OB, Anesthesia, Electrophysiology
- **Labor Management:**
 - **DPE placed at L4–5 + Prophylactic pacemaker pads**
 - **Initial epidural bolus:** 8 mL of 0.0625% bupivacaine (programmed intermittent)
 - Developed **HR 30 bpm**, BP 97/45 mmHg + dizziness
 - **Epidural paused**, T10 blockade confirmed
 - Treated with **10 mg IV ephedrine, 10 µg IV epinephrine** + Crystalloid bolus → Rapid recovery
 - Epidural restarted at **6 mL q30min** (down from 8 mL)
- **Outcome:**
 - No further arrhythmic events
 - **Uneventful vaginal delivery 9 hours later**
 - Epidural bolus of 100 mcg fentanyl and 5 cc 0.125% bupivacaine 2 hours prior to delivery without issue
 - Postpartum EP study → conduction lesion near AV node but **adequate chronotropic response** → No permanent pacemaker needed



Teaching Points:

- **Asymptomatic CCHB** still poses **peripartum risk**
- **Neuraxial anesthesia** → Sympathetic tone reduction can provoke bradycardia/asystole
- **Baseline HR <60** → Increased risk of moderate/severe bradycardia
- **Management Strategies:**
 - **Early labor induction** for worsening conduction
 - Prophylactic **temporary pacing pads**
 - **Titrated local anesthetic** dosing
 - **Immediate access** to vasopressors/inotropes (e.g., ephedrine, epinephrine)
 - **Multidisciplinary coordination** (OB, Anesthesia, Cardiology/EP)
- **Considerations for future patients with CCHB**
 - **Proactive Holter monitoring** before term to assess chronotropic competence
 - **Ensure immediate access to transvenous pacing**, especially in low-resource settings
 - **Engage EP/Cardiology early** in pregnancy for pacing strategy and follow-up

